



Review

Ketamine, magnesium and major depression – From pharmacology to pathophysiology and back

Harald Murck^{a,b,*}^a Covance Inc., Princeton, USA^b Clinic of Psychiatry and Psychotherapy of the Philipps-University of Marburg, Marburg, Germany

ARTICLE INFO

Article history:

Received 19 October 2012

Received in revised form

14 February 2013

Accepted 26 February 2013

Keywords:

Major depression

Ketamine

Magnesium

Synaptogenesis

NMDA

NR2b

CP-AMPA

mTOR

CaMKII

P2X7

Aldosterone

Slow wave sleep

Stress

ABSTRACT

The glutamatergic mechanism of antidepressant treatments is now in the center of research to overcome the limitations of monoamine-based approaches. There are several unresolved issues. For the action of the model compound, ketamine, NMDA-receptor block, AMPA-receptor activation and BDNF release appear to be involved in a mechanism, which leads to synaptic sprouting and strengthened synaptic connections. The link to the pathophysiology of depression is not clear. An overlooked connection is the role of magnesium, which acts as physiological NMDA-receptor antagonist: 1. There is overlap between the actions of ketamine with that of high doses of magnesium in animal models, finally leading to synaptic sprouting. 2. Magnesium and ketamine lead to synaptic strengthening, as measured by an increase in slow wave sleep in humans. 3. Pathophysiological mechanisms, which have been identified as risk factors for depression, lead to a reduction of (intracellular) magnesium. These are neuroendocrine changes (increased cortisol and aldosterone) and diabetes mellitus as well as Mg^{2+} deficiency. 4. Patients with therapy refractory depression appear to have lower CNS Mg^{2+} levels in comparison to health controls. 5. Experimental Mg^{2+} depletion leads to depression- and anxiety like behavior in animal models. 6. Ketamine, directly or indirectly via non-NMDA glutamate receptor activation, acts to increase brain Mg^{2+} levels. Similar effects have been observed with other classes of antidepressants. 7. Depressed patients with low Mg^{2+} levels tend to be therapy refractory. Accordingly, administration of Mg^{2+} either alone or in combination with standard antidepressants acts synergistically on depression like behavior in animal models.

Conclusion: On the basis of the potential pathophysiological role of Mg^{2+} -regulation, it may be possible to predict the action of ketamine and of related compounds based on Mg^{2+} levels. Furthermore, screening for compounds to increase neuronal Mg^{2+} concentration could be a promising instrument to identify new classes of antidepressants. Overall, any discussion of the glutamatergic system in affective disorders should consider the role of Mg^{2+} .

© 2013 Elsevier Ltd. All rights reserved.

1. Introduction

An interest in glutamatergic aspects in depression existed more than 20 years ago and important data were generated (Trullas and Skolnick, 1990; Paul et al., 1994). The renewed awareness of the limitations to treat depression with monoamine-based mechanism of action and the findings of a rapid antidepressant response to ketamine in patients with treatment resistance to these

compounds has led to a paradigm shift for the development of new antidepressant compounds. First observations of the effects of ketamine in depression (Berman et al., 2000) were done more than 10 years ago. However, it was not until the mile-stone observations of Zarate et al. (2006), who demonstrated in a double blind placebo controlled study the effect of ketamine in patients, who did not respond to standard antidepressants, that this field got broader attention. During the last several years has the NMDA ergic system received marked attention in the context of developing new compounds for mood disorder (Li et al., 2011b). The mechanism of action of glutamatergic compounds is unraveling, but there are several unresolved issues.

* Covance Inc., Princeton, USA. Tel.: +1 201 294 1195.

E-mail addresses: haraldmurck@yahoo.de, harald.murck@covance.com.

2. Proposed mechanism of action of ketamine in depression

The recently proposed sequence of events involves blocking N-methyl-D-aspartate (NMDA) receptors on gamma-aminobutyric acid (GABA)-ergic interneurons. The dampening of their activity disinhibits glutamatergic neurons due to a lowered GABAergic inhibition. Within the given context of NMDA blockade the released glutamate primarily excites (-amino-3-(3-hydroxy-5-methyl-iso-oxazol-4-yl) propanoic acid) (AMPA) and kainite receptors (Fig. 1). It is important to keep in mind that this combination of NMDA-receptor blockade and AMPAergic activation is of short duration due to the short half life of ketamine, but could be necessary for the long term antidepressant effects of ketamine treatment. 2. AMPA receptors activate the expression and release of brain derived neurotrophic factor (BDNF), which activate its receptor, the tropomyosin related kinase B (TrkB)-receptor.

Several alternative pathways downstream of the activation of TrkB receptors by BDNF have been described: a) activation of PI3K, followed by that of AKT and of mammalian target of rapamycin (mTOR) which leads to the inhibition of eukaryotic elongation factor 2 (eEF2) phosphorylation. Recently an alternative pathway via deactivation of eEF2-kinase has been reported. This pathway may not involve mTOR, but results in a reduced phosphorylation of eEF2 and disinhibition of BDNF translation (Autry et al., 2011). For both pathways, reduced phosphorylation of eEF2 and downstream disinhibition of gene expression seem to be relevant steps. b) A parallel pathway involves activation of extracellular signal regulated kinase (ERK). Via both pathways TrkB activation finally leads to an induction of synaptic proteins, in particular AMPAergic glutamate receptor subunits GluR1, anchoring protein PSD-95 and synapsin 1 (Li et al., 2010; Duman et al., 2012). The outcome of the BDNF related activation is therefore the increase in a specific subset of AMPA receptors (AMPA), i.e. Ca²⁺-permeable (CP)-AMPA (Fortin et al., 2012), when GluR1-subunits form homomeric receptors. As a note, other AMPAR, i.e. those which contain the GluR2 subunit, are impermeable for Ca²⁺ and Mg²⁺ (Burnashev et al., 1992). There is evidence that CP-AMPA are required for long term potentiation (LTP)-induced neuronal spine enlargement (Fortin et al., 2010), i.e. their increase may have specific relevance for synaptic sprouting. c) Finally, BDNF activates Ca²⁺/calmodulin-dependent protein kinase (CaM kinase) kinase (CaMKK) in a transient receptor potential canonical channel (TRPC)-dependent way, which activates AKT and in parallel mediates an increased synaptic incorporation of CP-AMPA (Fortin et al., 2012) (Fig. 2). It is not yet clear if this interaction between AMPA-receptor mediated BDNF release and BDNF related CP-AMPA expression constitutes a feedforward cycle, as it is not clear if a specific AMPAR subtype is involved in the BDNF expression and release.

Opposing mechanisms for the therapeutically relevant switch of AMPA-receptor constitution have to be considered: Extrasynaptic NMDA-receptor activation, which occurs during glutamate spillover in pathogenetic processes leads to an inhibition of CREB and reduces BDNF expression (Vanhoutte and Bading, 2003). This leads to a PKC dependent switch from CP-AMPA to GluR2 containing Ca²⁺ and Mg²⁺ impermeable channels (Sun and June Liu, 2007). This mechanism may in part explain the action of compounds like riluzole. Riluzole reduces the extracellular concentration of glutamate and may therefore reduce glutamate spillover to extrasynaptic NMDA receptors (Sanacora et al., 2004, 2007). Therefore, there is a tight window of a beneficial concentration of synaptic glutamate, which primarily acts on synaptic AMPAR and a potentially deleterious spillover to extrasynaptic NMDA receptors. The combined AMPAR activation and NMDA-receptor inhibition at the early stages of the effect of ketamine may therefore be of importance for ketamine's clinical effect.

Several questions occur: What determines the response to ketamine in a given subject? What determines the release of BDNF? Is the release of BDNF with ketamine in any way different than that with standard antidepressants?

3. Comparison of the effects of Mg²⁺ and ketamine

The comparison of the effect of ketamine with the physiologically present NMDA antagonistic Mg²⁺ may provide further insights into the mechanisms in question. 1. Ketamine- and Mg²⁺ administration both lead to an increase in slow wave sleep in humans preferably at the beginning of the sleep period (Held et al., 2002; Duncan et al., 2012b). 2. Administration of an innovative Mg²⁺ compound with a high bioavailability leads to an increase in BDNF expression in the brain and finally to synaptic sprouting in the prefrontal cortex of rats, again, similar to ketamine (Abumaria et al., 2011). In parallel an increase in NR2b (but not NR1 and NR2a) receptor expression occurred with both ketamine (Burgdorf et al., 2013; Chatterjee et al., 2012) and Mg²⁺ – administration (Abumaria et al., 2011). Opposite changes can be observed in neuronal cell cultures exposed to short term Mg²⁺-free solution, which leads to a reduced expression of NR2b and PSD-95 (Jiang et al., 2007). 3. Chronic unpredictable stress reduces neuronal strength, i.e. the amplitude of excitatory postsynaptic potential (EPSP) in the PFC of rats, which can be reversed by the administration with ketamine (Li et al., 2011a). Similarly, incubation of prefrontal cortex slices with Mg²⁺ increase of EPSP amplitude (Abumaria et al., 2011). It would be of interest to determine, if incubation with Mg²⁺ leads to a normalization of the stress-induced EPSP reduction. Overall, the functional consequences of Mg²⁺ and ketamine administration show similarities.

4. Overlap of Mg²⁺ and ketamine-induced pathways

4.1. NR2b antagonistic effect of Mg²⁺

Mg²⁺ is a naturally occurring NMDA-receptor antagonist and has effects in concentrations, which are physiologically occurring in the extrasynaptic space (see Murck, 2002). Of particular interest is the relative specificity of the Mg²⁺ block of the NMDA receptors based on their specific subunits. The NR1 receptor is ubiquitous, whereas different kinds of NR2 subunits exist. Of these the NR2c and NR2d are the least sensitive to the Mg²⁺ block, whereas NR2a and NR2b are more sensitive (Kuner and Schoepfer, 1996). It is of interest that it is the NR2b receptor, which is a target for antidepressive efficacy similar to that of ketamine (Preskorn et al., 2008). Is therefore ketamine potentially replacing a deficit of Mg²⁺ at the level of the NR2b type NMDA receptor? There are a number of observations, which appear to be in line with this notion: the sensitivity of ketamine and other NMDA-receptor antagonists is increased in a state of Mg²⁺ depletion (Begon et al., 2001). In particular Mg²⁺ deficiency leads to a sensitization for ketamine to induce sleep (Douglas and Dagirmanjian, 1975). In line with this is that treatment refractory depressed patients, i.e. the group which shows beneficial effects from ketamine treatment, have a lower CNS Mg²⁺ level (Iosifescu et al., 2008). However, this may not be the complete explanation. There are some observations, which demonstrate that the combination of ketamine and Mg²⁺ in a situation of normal Mg²⁺ levels has super-additive effects (Orser et al., 1997; Liu et al., 2001). Therefore potentially synergistic effects beyond NR2b antagonism should be taken into account.

This NR2b antagonistic effect of Mg²⁺ is complemented by intracellular mechanism of Mg²⁺, for an overview see (Murck, 2002). It has to be considered that Mg²⁺ is a primarily intracellular ion. Therefore the reduced Mg²⁺ level in the brain of subjects




Fig. 1. Magnesium involvement in ketamine-induced pathways: ketamine administration leads to a cascade of events finally resulting in modifications of glutamatergic receptor profile and synaptogenesis. Functional consequences are increased excitatory postsynaptic potentials (EPSP) and increased slow wave sleep; both phenomena are also induced by Mg²⁺. In detail: A: ketamine inhibits GABAergic interneurons and therefore activates the release of glutamate. B: AMPAR mediate BDNF expression and release. BDNF activates TrkB receptor, which induces changes in gene expression (see Fig. 2). BDNF induces its own expression. NMDA-receptor activation is facilitating this process (Xiong et al., 2002). This could constitute a feedforward mechanism, explaining the long term effect of ketamine administration. Further, the expression of synaptic proteins is induced, in particular GluR1 and PSD-95, which constitute the synaptic expression of Ca²⁺-permeable AMPA receptors (CP-AMPA) (Fortin et al., 2012). Importantly, CP-AMPA are permeable for Mg²⁺. C: Glutamate is taken up quickly by neurons and astrocytes. This is of importance as high concentrations of glutamate can “spillover” to extrasynaptic NMDA receptors, which appear to be primarily from the NR2b type. Their activation can block synaptogenesis and lead to cell damage. Rapid reuptake of glutamate prevents this “spillover”. Glutamate reuptake is an energy dependent process driven by the Na⁺-gradient over the membrane (Magistretti, 2009), which in itself is driven by the Na⁺-K⁺-ATPase. The Na⁺-K⁺-ATPase is dependent on Mg²⁺, therefore increased Mg²⁺ availability supports its activity and secondarily glutamate clearance. D: In parallel glutamate receptors at astrocytes are activated. CP-AMPA receptor activation can lead to an increase in astrocytic Ca²⁺ (Magistretti, 2009) and Mg²⁺ (Muller et al., 2003). A potential consequence of increased Mg²⁺ in astrocytes is activation of glutamine synthetase (GS), which is Mg²⁺ dependent (Greenberg and Lichtenstein, 1959; Maurizi et al., 1986). E: A neuronal Na⁺-Mg²⁺-exchange mechanism regulates intracellular Mg²⁺ concentration. Imipramine appears to block Na⁺-Mg²⁺-exchange, preventing the efflux of Mg²⁺ from the neuron. F: Magnesium uptake into the brain has been described with compounds, which are known to be efficacious in treatment refractory depression, i.e. ketamine (via its glutamate releasing capability), TSH, lithium, imipramine and potentially insulin related mechanism (metformin, glitazones). On the other hand stress and refractory depression are linked to lower Mg²⁺ levels in the brain, which may in part be mediated via an aldosterone mediated mechanism.

with depression should be reflected in intracellular changes as well. The intracellular Mg²⁺ concentration affects the sensitivity of NMDA receptors and their activity: In sensory neurons (Chen and Huang, 1992) and in hippocampal synaptosomes, the Mg²⁺ block of the NMDA dependent ion channel is removed by activation of protein kinase C (PKC) without changing membrane potential (Pittaluga et al., 2000). Intracellular administration of a PKC agonist accordingly potentiated NMDA-receptor function in cultured hippocampal neurons (Xiong et al., 1998). On the other hand Mg²⁺ appears to affect PKC activity: Mg²⁺ depletion leads to an increase in NMDA mediated PKC activation and nitric oxide (NO) release in spinal cord neurons (Begon et al., 2001). This mechanism could lead to a feedforward cycle: a NMDA dependent Ca²⁺ current may increase PKC activity, which leads to further release of the Mg²⁺ block of the NMDA dependent ion current (Chen and Huang, 1992). This means that reduced intracellular Mg²⁺ can be responsible for an increased NMDA-receptor sensitivity. An alternative mechanism of intracellular Mg²⁺ deficiency to increase NR2b-receptor activation exists, which appears to be independent from second messenger pathways (Li-Smerin et al., 2000). Overall, both extracellular and intracellular Mg²⁺ inhibit NMDA-receptor activity and in particular NR2b-receptor function.

4.2. Comparison of the effects of ketamine and Mg²⁺ on the CaMKII pathway

An increase in synaptic strength appears to be the result of the effects of both ketamine and Mg²⁺. One of the molecular determinants for synaptic strength is the interaction between NR2b containing NMDA receptors on the one hand and Ca²⁺/calmodulin activated protein kinase II (CaMKII) and PSD-95 proteins (Lisman et al., 2012). Increase in intracellular Ca²⁺ leads to the activation of calmodulin, which activates CaMKII. This leads to a translocation and fixation of GluR1 containing AMPA receptors at synaptic sites and as a consequence amplification of AMPAergic currents (Lee et al., 2000; Lisman et al., 2012). This mechanism increases EPSPs frequency and long term potentiation (LTP) in the hippocampus.

Mg²⁺ leads to increased LTP in hippocampal slices (Landfield and Morgan, 1984). Low Mg²⁺ for a short period of time leads to an inhibition of CaMKII activity (Blair et al., 1999), whereas Mg²⁺ administration in vivo increases phosphorylation and therefore activation of CaMKII in the prefrontal cortex. This was accompanied by an increase in LTP (Abumaria et al., 2011). Similarly ketamine induces CaMKII expression, as observed in the nucleus accumbens of rats (Iasevoli et al., 2007). Interestingly a reduced expression of




Fig. 2. Proposed pathways of synaptogenesis. A. BDNF activated TrkB receptors lead to the activation of several parallel pathways including: The activity of a) the AKT–mTOR pathway and as a consequence inhibition of eEF2-kinase. The eEF2-kinase appears to be directly inhibited by Mg²⁺ (Perraud et al., 2011). b) The ERK pathway, which is also facilitated by Mg²⁺ administration (Huang et al., 2010); c) TPRC dependent activation of CaMKK and activation of AKT (Fortin et al., 2012). Gene expression is activated to lead to the generation of synaptic proteins, in particular GluR1 containing CP-AMPA as well as PSD-95. In parallel CaMKK activation is involved in the fixation of AMPAR at synaptic sites. B. The insertion of CP-AMPA is additionally regulated via CaMKII activation via NMDA-receptor related Ca²⁺ increase (Barria and Malinow, 2005). The CaMKII pathway is activated by Mg²⁺ and inhibited by P2X7-purinergic receptors, which by themselves are sensitive to Mg²⁺. As a consequence Mg²⁺ facilitates CaMKII-dependent synaptic activity. The resulting increased availability of CP-AMPA may lead to a higher sensitivity of the system to Mg²⁺, forming a feedforward cycle. C. The activation of extrasynaptic NMDA receptors by glutamate spillover appear to alter the AMPA-receptor expression by favoring GluR2 containing AMPAR (Non CP-AMPA) in a PKC dependent pathway (Sun and June Liu, 2007). This can have deleterious effects.

CaMKII in post mortem studies of the prefrontal cortex of patients with unipolar depression and bipolar disorder (Xing et al., 2002) has been observed, in line with the hypothesis of an involvement of Mg²⁺ dysregulation in patients with depression. In summary, Mg²⁺ increases CaMKII activity, which leads to an increase in AMPAergic function in the prefrontal cortex (Fig. 2).

In addition to the direct effect of Mg²⁺ on CaMKII activity an indirect pathway of importance exist. We want to focus here on the role of P2X7, a purinergic receptor, which controls an unspecific cation current (Skaper et al., 2010). Polymorphisms of the P2X7 gene have been linked to the risk of depression (Lucae et al., 2006; Soronen et al., 2011), however a negative study also exists (Green et al., 2009). In our context it is important that P2X7 receptor inhibition activates neurogenesis in a CaMKII-dependent pathway (Leon et al., 2006; Gomez-Villafuertes et al., 2009) in neuronal cells. As Mg²⁺ demonstrates an antagonistic effect on P2X7 receptor activation, as demonstrated in a variety of cells (Jiang, 2009; Alloisio et al., 2010; Lee et al., 2011) (Fig. 2), this points to a beneficial effect of Mg²⁺ in increase synaptic strength by reducing P2X7 activity.

4.3. cAMP–CREB–BDNF pathway

A further observation after Mg²⁺ administration was the increased phosphorylation of CREB (Abumaria et al., 2011). The importance of CREB-phosphorylation as a mediator of an antidepressant response has been extensively described (D'Sa and Duman, 2002; Carlezon et al., 2005; Gourley et al., 2008; Tanis et al., 2008). The supportive effect of Mg²⁺ on the activation of the cAMP pathway is well documented (see (Murck, 2002)). A direct importance of CREB for BDNF expression has been demonstrated (Duman, 1998; Manji and Duman, 2001). As a consequence, Mg²⁺ administration led to an increase in BDNF expression (Abumaria et al., 2011) and shows neuroprotective effects via CREB-phosphorylation (Huang et al., 2010).

4.4. AKT–mTOR–eEF2k pathway

The AKT–mTOR–eEF2k pathway activation appears to be one prominent consequence of TrkB activation. Mg²⁺, like ketamine, suppresses eEF2 phosphorylation (Perraud et al., 2011) in cell

Table 1

Overview of discussed changes in regional expression of receptors and transcription factors if not otherwise specified the results are from animal models.

	Major depression (human)	Chronic stress	Mg depletion (acute)	Magnesium administration	Ketamine	Sleep deprivation
PFC–NR1	⇔ (Feyissa et al., 2009)	↓ (Lee and Goto, 2011)	?	⇔ (Abumaria et al., 2011)	↑ (Chatterjee et al., 2012)	?
Amygdala–NR1	⇔ (Karolewicz et al., 2009)	?	?	⇔ (Abumaria et al., 2011)	?	?
PFC–NR2A	↓ (Feyissa et al., 2009)	↓ (Lee and Goto, 2011; Yuen et al., 2012)	?	⇔ (Abumaria et al., 2011)	⇔ (Chatterjee et al., 2012)	8 h: ↑ (Cirelli and Tononi, 2000); 72 h: ⇔ (Park et al., 2012; Vyazovskiy et al., 2008)
Amygdala–NR2A	↑ (Karolewicz et al., 2009)	?	?	⇔ (Abumaria et al., 2011)	?	?
PFC–NR2B	↓ (Feyissa et al., 2009)	↓ (Lee and Goto, 2011; Yuen et al., 2012)	↓ (Jiang et al., 2007) (cell culture)	↑ (Abumaria et al., 2011)	↑ (Chatterjee et al., 2012; Burgdorf et al., 2013)	72 h: ↓ (Park et al., 2012)
Cortical (PFC)–GluR1	?	↓ (Li et al., 2011a; Yuen et al., 2012)	↑ (Jiang et al., 2008) (cell culture)	?	↑ (Li et al., 2011a)	3 h: ↑ (Vyazovskiy et al., 2008; after wake: Lante et al., 2011)
Cortical (PFC)–GluR2	↑ (Teyssier et al., 2011)	↓ (Yuen et al., 2012)	Biphasic: ↑ → ↓ (Jiang et al., 2008) (cell culture)	?	?	8 h: ↑ (Cirelli and Tononi, 2000); 3 h: ⇔ (Vyazovskiy et al., 2008)

Table 2

Overview of discussed changes in regional expression of transcription factors if not otherwise specified the results are from animal models.

	Major depression (human)	Chronic stress	Mg depletion (acute)	Magnesium administration	Ketamine	Sleep deprivation
PFC–BDNF	?	⇔ (Chiba et al., 2012); ↓ (van Donkelaar et al., 2009; Yu et al., 2011)	?	↑ (Abumaria et al., 2011)	↑ (Reus et al., 2011)	8 h: ↑ (Cirelli and Tononi, 2000; 3 h: Vyazovskiy et al., 2008); 72 h: ↓ (Park et al., 2012)
Amygdala–BDNF	?	⇔ (Reus et al., 2011); ↑ (Yu et al., 2011)	?	⇔ (Abumaria et al., 2011)	↑ (Reus et al., 2011)	?
PFC–PSD-95	↓ (Feyissa et al., 2009)	↓ (Li et al., 2011a; Zhang et al., 2012)	↓ (Jiang et al., 2007) (cell culture)	?	↑ (Li et al., 2011a) (after CUS)	?
Amygdala–PSD-95	↑ (Karolewicz et al., 2009)	↑ (Zhang et al., 2012)	?	?	?	?
Cortical P-CREB	↓ (Yamada et al., 2003; Dwivedi et al., 2003)	↓ (Kuipers et al., 2003; Laifenfeld et al., 2005)	?	↑ (Huang et al., 2010); ⇔ (Abumaria et al., 2011)	↑ (Reus et al., 2011; Shu et al., 2012)	8 h or 48 h: ⇔ (Guzman-Marin et al., 2006)
PFC/ACC–glutamine/GS activity	↓ (Choudary et al., 2005)	↓ (Knox et al., 2010; Hemanth Kumar et al., 2012)	?	↑ (Maurizi et al., 1986)	↑ (Rowland et al., 2005)	Human MDE: ↑ (Murck et al., 2009; 12–14 h: Bettendorff et al., 1996)
Evoked synaptic potentials (cortex)	?	↓ (Quan et al., 2011; Yuen et al., 2012)	↓ (Richards and Sercombe, 1970 ^a)	↑ (Abumaria et al., 2011)	↑ (Li et al., 2011a) (after CUS)	3 h: ↑ (Vyazovskiy et al., 2008)
Cortical synaptogenesis	?	↓ (Li et al., 2011a)	?	↑ (Abumaria et al., 2011)	↑ (Li et al., 2011a) (after CUS)	↑ ((Maret et al., 2011)

^a Supraphysiological concentrations reduced EPSP amplitude.

culture and increased NR2b receptor and BDNF expression in the hippocampus, increases synaptic connectivity and synaptogenesis (Slutsky et al., 2010). This may point to a parallel mechanism of Mg^{2+} to amplify the BDNF induced cascade.

In summary, Mg^{2+} is involved in a number of mechanisms, which have central relevance for the pathophysiology of major depression and which are also affected by ketamine administration. The issue arises if some of the effects of ketamine are related to changes in intracellular Mg^{2+} .

5. Importance of regional differentiation

The regional differentiation of the biological effects of stress and antidepressant treatments is of critical importance (Tables 1 and 2): Opposite change of BDNF expression in the PFC vs. the amygdala occurs under stressful conditions (Yu and Chen, 2011). A similar regional specificity of the effects of ketamine and Mg^{2+} administration exists: Mg^{2+} administration leads to an increase in NR2b-receptor expression, as well as to CaMKII-phosphorylation, CREB-phosphorylation and BDNF expression, but not that of NR1, in the prefrontal cortex, but not in the amygdala of rats (Abumaria et al., 2011). Similarly, ketamine leads to an increase in CREB-expression and PKC phosphorylation in the prefrontal cortex, but not in the amygdala (Reus et al., 2011). BDNF is increased after both, ketamine and Mg^{2+} administration in the prefrontal cortex. Small differences exist: ketamine increases BDNF in the amygdala as well, whereas Mg^{2+} does not have an effect in the tested dose.

A similar regional specificity of the gene expression is observed in patients with depression (Tables 1 and 2). In post mortem studies of brains of patients with depression reduced levels of NR2a and NR2b subunits have been identified, accompanied by a reduction of PDS-95 (Beneyto and Meador-Woodruff, 2008; Feyissa et al., 2009). In the lateral amygdala partially opposite changes were observed, i.e. elevated levels of NR2a and PDS-95 (Karolewicz et al., 2009). Therefore the changes in the prefrontal cortex are the mirror image of what was observed with Mg^{2+} administration. The changes in the amygdala differ from those in the prefrontal cortex in both instances (Tables 1 and 2).

The relationship between major depression and chronic stress is well established and is reflected in changes of glutamatergic receptor expression (Tables 1 and 2). One finding should be specifically highlighted, which has relevance for the further argumentation: the prefrontal cortex in patients with depression shows an increased expression of GluR2 type subunit of the AMPA receptor, i.e. non CP-AMPA (Teyssier et al., 2011). Furthermore stress in an animal model increases GluR2-phosphorylation and therefore their recruitment to postsynaptic sites in the PFC, but has opposite effects in the amygdala (Caudal et al., 2010). This highlights the importance of regional specificity of AMPAR constitution in stress related situations (Tables 1 and 2).

6. Glutamatergic regulation of CNS Mg^{2+} content

It appears that the mechanism of action of ketamine and Mg^{2+} overlaps beyond the NMDA antagonistic effect of extracellular Mg^{2+} . The question is, how may this happen? An interesting observation is that ketamine actually leads to an increase of intracellular Mg^{2+} in peripheral tissue (Kim et al., 2006), which involves an increase in ERK1/2 and p38 MAP kinase. Further ketamine and MK-801 reverses the decrease of brain Mg^{2+} after brain trauma (McIntosh et al., 1990; Shapira et al., 1993). The described antidepressant mode of action of ketamine involves an increase of glutamate the prefrontal cortex and an activation of non-NMDA receptors. I.c.v. administered kainite and quinolinate leads to an increase in intracerebral Mg^{2+} content (against a concentration

gradient from the plasma) (Rothe et al., 1993). At the cellular level glutamate leads to an increase in intracellular Mg^{2+} concentration in cultured rat forebrain neurons (Cheng and Reynolds, 2000), however in very artificial conditions (Na^+ and Ca^{2+} free medium). This increase consists of two elements; firstly an increase of intracellular Mg^{2+} -release and secondly an influx of Mg^{2+} in a Na^+ dependent way (Hoyt et al., 1995). Non-NMDA glutamate receptors (AMPA/kainite receptors) mediate this increase in intracellular Mg^{2+} (Hoyt et al., 1995). As we saw earlier the effect of ketamine at the postsynaptic levels is to increase the expression of CP-AMPA. As CP-AMPA are permeable to Mg^{2+} , this could be a potential mechanism to increase intracellular Mg^{2+} levels as a consequence of ketamine action. Therefore one mechanism of action of ketamine could be to increase intracerebral Mg^{2+} content via glutamatergic stimulation.

7. Synoptic overview of rapid antidepressant interventions

As an overview of the discussed mechanism a comparison of the effects of the different manipulations is provided in Tables 1 and 2. Additional information on the effects of chronic stress, which can be regarded as a model of depression, is included (Kuipers et al., 2003; Laifenfeld et al., 2005; van Donkelaar et al., 2009; Knox et al., 2010; Lee and Goto, 2011; Quan et al., 2011; Yu and Chen, 2011; Chiba et al., 2012; Hemanth Kumar et al., 2012; Yuen et al., 2012; Zhang et al., 2012) as subchronic stress related changes are reversed by ketamine in an animal model (Li et al., 2011a). Table 1 provides a synopsis of the described conditions at the level of glutamate receptors, whereas Table 2 provides an overview of downstream changes. From this overview it becomes plausible that the changes in chronic stress (in animal models) resemble those in patients with major depression. This includes glutamate receptor composition, but also intracellular signal mechanism, like CREB-phosphorylation (Dwivedi et al., 2003; Yamada et al., 2003). One noteworthy exception is the regulation of the GluR2 AMPA-receptor component. As his parameter depends on the status of previous sleep variability in human post mortem material is expected. There is some overlap between the changes seen in Mg^{2+} depletion and chronic stress, however, the Mg^{2+} -depletion data are mainly from cell culture experiments. One study observed electrophysiological effects of several mainly supraphysiological Mg^{2+} -concentrations on evoked potentials in brain slices (Richards and Sercombe, 1970). Gene expression in animal models with Mg^{2+} depletion has not been studied in the areas of interest for the receptors of interest. Mg^{2+} and ketamine administration generally has the opposite effect than those observed in chronic stress models. An important and potentially clinically relevant difference is that ketamine leads to an increase in amygdala-BDNF, which may be related to its potential to induce unpleasant phenomena, including psychotic symptoms.

The action of ketamine has similarities with that of therapeutic sleep deprivation (TSD) (Zarate et al., 2013). Sleep deprivation, has opposite effects on receptor expression than chronic stress, however this is only the case for short term (up to 24 h) sleep deprivation. Longer term sleep deprivation has partially opposite effects. Both TSD and ketamine are fast acting; they lead to similar polysomnographic changes, i.e. primarily an increase in slow wave sleep; both increase glutamine levels in the prefrontal cortex (Rowland et al., 2005; Murck et al., 2009; Duncan et al., 2012b; Taylor et al., 2012). Glutamine synthetase (GS) activity is increased in the prefrontal cortex in animal models after sleep deprivation (Bettendorff et al., 1996). Therefore a consistent effect of the antidepressant interventions described is the increase in GS activity, i.e. of the enzyme which catalyzes glutamine synthesis. Further GS has been related to suicide and depression (Choudary et al., 2005;

Kalkman, 2011). The increase in glutamine is correlated with the reduction in depressive mood after therapeutic sleep deprivation (Murck et al., 2009) and potentially ketamine administration (Salvadore et al., 2011). AMPAR activation (Fleischer-Lambropoulos et al., 1996) followed by intracellular Mg^{2+} increase could be responsible for this, as GS is a Mg^{2+} dependent enzyme (Greenberg and Lichtenstein, 1959; Maurizi et al., 1987). In summary, administration of ketamine mimics the effect of sleep deprivation in a molecular, electrophysiological and clinical level. For a wider discussion on the mechanism of sleep deprivation see (Hemmeter et al., 2010).

8. Predictors of ketamine response

It is important to note that not all patients with depression can be expected to respond to ketamine. Studies have been done in patients with therapy refractory depression, who may represent a specific subtype of all depressed patients. In this population the number of patients needed to treat (NNT) is 3–5, which is an extremely good effect, however not complete (Aan Het Rot et al., 2012). Predictors of response for ketamine are being studied: Reduced SWS at the beginning of the sleep period, as expressed as delta sleep ratio, is a positive predictor for the effect of ketamine (Duncan et al., 2012a). It is the opposite of the effect of Mg^{2+} (Held et al., 2002). Furthermore the increase in slow wave sleep is correlated to the clinical response in responders to ketamine (Duncan et al., 2012b). Interestingly, in mice frontal cortical Mg^{2+} concentration is correlated with the increase in slow wave sleep after short term sleep deprivation (Chollet et al., 2000). A similar correlation was found with red blood cell Mg^{2+} in the same study. In addition, lower dorsomedial PFC glutamine levels predict a preferable outcome of the ketamine effect on depressive symptoms (Salvadore et al., 2011). Given that GS is a Mg^{2+} dependent enzyme, both of these findings are in line with a potential role of low (intracellular) Mg^{2+} content as a predictor of ketamine response.

Direct evidence for a reduction of intracerebral Mg^{2+} in this particular group of patients with therapy refractory depression comes from a study utilizing Phospho-(P)-spectroscopy (Iosifescu et al., 2008). Mg^{2+} is primarily located intracellularly, therefore reduced overall brain Mg^{2+} level points toward a reduction of its intracellular content. Similarly, the Mg^{2+} content of red blood cells is reduced in patients with depression (Nechifor, 2008). For a recent overview see Eby et al. (2011). Furthermore, subjects with lower levels of Mg^{2+} , as measured in plasma, appear to have worse response to standard antidepressant treatment (Camardese et al., 2012). In conclusion, markers of low intracellular Mg^{2+} content may predict preferable response to ketamine. The most direct way to measure intracellular Mg^{2+} is by means of P-spectroscopy. Other direct measures of intracellular Mg^{2+} may utilize red blood cells (Nechifor, 2008), white blood cells (Ahokas et al., 2003) or epithelial cells (Silver, 2004). A potential functional marker may be the amount of SWS or delta power of the sleep EEG at the beginning of the night.

9. Pathophysiological connections

If some forms of depression may be mediated via low intracerebral Mg^{2+} and ketamine potentially reverses these changes, the question is, what led to these low Mg^{2+} level? Is there a connection to established mechanism of depression? In fact, Mg^{2+} is excreted by an increased activity of the sympathetic nervous system and the hypothalamus–pituitary adrenocortical axis (Murck, 2002; Murck et al., 2012). It has been suggested that the pathophysiological changes targeted by ketamine originate from increased cortisol or corticosterone mediated glucocorticoid receptor (GR) activation

(Popoli et al., 2012). A recent observation, however, points to the potential involvement of mineralocorticoid receptors (MR). Corticosterone induced depression like behavior was counteracted by the administration of the MR antagonist spironolactone (Wu et al., 2012). Therefore the physiological MR agonistic corticosteroid, i.e. aldosterone, should be taken into account: hyperaldosteronism has been reported in major depression (Murck et al., 2003; Emanuele et al., 2005). Aldosterone induces depressive symptoms in both animal models (Hlavacova et al., 2011) and humans (Kunzel et al., 2012). On the basis of the timing of neuroendocrine changes a causal role of aldosterone in diverse animal models has recently been suggested (Franklin et al., 2012). Given that Mg^{2+} excretion is activated by aldosterone it makes therefore physiological sense that Mg^{2+} could be the link between stress related behavioral changes and overactivity of the NMDA ergic system.

10. Increase in intracellular Mg^{2+} as a mediator for antidepressant efficacy

I summarized the parallel effects between ketamine and Mg^{2+} . I also provided evidence that Mg^{2+} is involved in pathways, which are connected to the known pathophysiology of depression. Further I described mechanisms, which lead to an increase in intracerebral intracellular Mg^{2+} . The question is now, is there precedence for the action of standard antidepressants on Mg^{2+} ? There are several observations, which point into that direction: Of high interest is that imipramine, the first reuptake inhibitor and therefore the model substance for most currently used antidepressants, leads to an increase in intracellular Mg^{2+} concentration in peripheral tissue and the brain (Poleszak et al., 2005; Lee et al., 2010). Just consider that this observation would have been done before the monoamine depletion hypothesis had been generated and the pharmacology of depression would probably look very different today. The mechanism of action for imipramine in erythrocytes is to inhibit a Na^+ – Mg^{2+} -exchange transporter at physiological intracellular Mg^{2+} levels (Ebel et al., 2004) and therefore to increase intracellular Mg^{2+} , i.e. to reduce the Mg^{2+} -transport out of the cell. An increase in intracerebral Mg^{2+} levels has also been demonstrated with compounds, which are recommended as adjunct therapy for therapy refractory depression: Thyroid stimulation hormone (TSH) has been demonstrated to increase brain Mg^{2+} content in patients with depression (Iosifescu et al., 2008). We already mentioned imipramine, which is still regarded as more efficacious than SSRIs. Finally, lithium increases Mg^{2+} in neuroblastoma cells (Amari et al., 1999; Abukhdeir et al., 2003) and in animal models after several days of treatment (King et al., 1969), but only by trend after 8 weeks of treatment (Kielczykowska et al., 2007). Therefore there is some indication that the effect of lithium on brain Mg^{2+} is time dependent and potentially has an inverted U-shaped characteristic, as high doses lead actually to a decrease (Kielczykowska et al., 2003). This finding is in line with the effect of lithium on erythrocyte Mg^{2+} , which shows acutely an increase, but no change in the long term, pointing to the importance of adaptive changes (Dunner et al., 1975).

Compounds with properties to increase intracellular Mg^{2+} express antidepressive efficacy in preliminary animal experiments or preliminary clinical trials: These are for example mineralocorticoid receptor antagonists (Ahokas et al., 2003; Runyan et al., 2005); which protect against aldosterone induced Mg^{2+} depletion and by this mechanism may have an antidepressant effect (Ebel et al., 1971; Hlavacova et al., 2011; Murck et al., 2012). Antidiabetic drugs of the glitazone group lead to an increase of Mg^{2+} in peripheral mononuclear cells in vitro (Alon et al., 2006) and have antidepressant effects in animal models (Salehi-Sadaghiani et al., 2012) and in clinical trials (Rasgon et al., 2010; Sepanjnia et al.,

2012). Metformin leads to an increase in intracellular Mg^{2+} levels in vivo (Gorelik et al., 2007), however its effect on depression is not clear (Rubin et al., 2005). These Mg^{2+} -increasing compounds act via different mechanisms of action, therefore the outcome, i.e. the increase in intracellular Mg^{2+} , could be regarded as a common read-out with relevance for the downstream effects, i.e. modification of glutamatergic mechanism as described above.

Following the suggestion that certain antidepressant mechanisms are related to an increase in Mg^{2+} it would be expected that there is a synergism between these compounds and Mg^{2+} administration. Indeed, it has been demonstrated that Mg^{2+} deficiency leads to depression like behavior in rats, which is reversed by imipramine (Singewald et al., 2004). Furthermore it has been demonstrated that the combination of sub-therapeutic doses of Mg^{2+} in combination with sub-therapeutic doses of imipramine leads to a significant antidepressive like effect in animal models (Poleszak et al., 2005, 2006). Further, the effect of the NMDA antagonist MK-801, which is similar to ketamine, can be amplified by concomitant administration of Mg^{2+} (Poleszak et al., 2007).

11. Conclusion

What are the conclusions of the described relationships? These are at least threefold:

1. Reduced concentration of Mg^{2+} in several tissues or functional markers of reduced tissue Mg^{2+} content appear to indicate worse response to antidepressant therapy with standard, the monoamine system targeting compounds. Direct intracellular measures in peripheral cells, P-spectroscopy of the brain or functional markers, like the pattern of slow wave sleep may therefore be predictive for the response of antidepressant compounds. Given the inverted U-shaped dose–response curve for the clinical ketamine response, functional markers like polysomnography, may help to define the individualized titration of compounds, which affect the glutamatergic system.
2. The property of compounds to increase intracellular Mg^{2+} levels could be a new biological target for antidepressants, primarily for the treatment of standard therapy refractory depression. This would open up a new strategy for candidate compound identification and selection.
3. A combination therapy may be considered, in particular with compounds, which increase intracellular Mg^{2+} and with Mg^{2+} itself. For example, the combination of NMDA-antagonism and Mg^{2+} has been demonstrated to show synergistic effects in animal models. Similarly, plasma Mg^{2+} levels appear to correlate to the outcome with standard antidepressant treatment (Camardese et al., 2012). Preliminary clinical observations point to the usefulness of concomitant administration of Mg^{2+} adjunct to antidepressive therapy (Eby and Eby, 2006; Barragan-Rodriguez et al., 2008), however, negative results after short term i.v. administration in patients with premenstrual syndrome also exist (Khine et al., 2006).

One point of caution: All the mechanism discussed focused on direct CNS effects. Of potential importance are immunological (Loix et al., 2011; Zeng et al., 2011) and neuroendocrine (Broughton Pipkin and Waldron, 1983; van Berckel et al., 1998) effects of ketamine, as well as of Mg^{2+} (Weglicki et al., 1992; Sartori et al., 2012; Weglicki, 2012). These peripheral mechanisms have been demonstrated to be involved in the pathophysiology of depression and should not be neglected.

As the final conclusion Mg^{2+} appears to be an important player in the pathophysiology of some forms of depression. Mg^{2+} related markers have relevance for treatment prediction with glutamatergic compounds and increased Mg^{2+} may moderate antidepressant mechanisms. Therefore, a discussion of glutamatergic mechanism of affective disorders should take the role of Mg^{2+} into account.

Conflict of interest

Harald Murck works currently for Covance Inc. Princeton, USA. He was formerly employed by Bristol-Myers Squibb, Novartis, Amarin and Lichtwer Pharma. There exists no conflict of interest. No funding for the preparation of this article was received.

Contributors

Harald Murck is the sole contributor to this article.

Role of the funding source

No funding was received for preparing this article.

Acknowledgment

I would like to thank Ronald Duman, Yale University and Carlos Zarate, NIMH, and Laura Zumpano, Pfizer, for their helpful comments.

References

- Aan Het Rot M, Zarate Jr CA, Charney DS, Mathew SJ. Ketamine for depression: where do we go from here? *Biological Psychiatry* 2012.
- Abukhdeir AM, Layden BT, Minadeo N, Bryant FB, Stubbs Jr EB, Mota de Freitas D. Effect of chronic Li^+ treatment on free intracellular Mg^{2+} in human neuroblastoma SH-SY5Y cells. *Bipolar Disorders* 2003;5:6–13.
- Abumaria N, Yin B, Zhang L, Li XY, Chen T, Descalzi G, et al. Effects of elevation of brain magnesium on fear conditioning, fear extinction, and synaptic plasticity in the infralimbic prefrontal cortex and lateral amygdala. *Journal of Neuroscience* 2011;31:14871–81.
- Ahokas RA, Warrington KJ, Gerling IC, Sun Y, Wodi LA, Herring PA, et al. Aldosterone and peripheral blood mononuclear cell activation: a neuroendocrine-immune interface. *Circulation Research* 2003;93:e124–35.
- Alloisio S, Di Garbo A, Barbieri R, Bozzo L, Ferroni S, Nobile M. Evidence for two conductive pathways in P2X receptor: differences in modulation and selectivity. *Journal of Neurochemistry* 2010;113:796–806.
- Alon I, Berman S, Shteinshnaider M, Efrati S, Gorelik O, Almozino-Sarafian D, et al. Total cellular calcium and magnesium content of cultured human peripheral blood mononuclear cells following exposure to antidiabetic drugs. *Acta Diabetologica* 2006;43:46–51.
- Amari L, Layden B, Nikolakopoulos J, Rong Q, Mota de Freitas D, Baltazar G, et al. Competition between Li^+ and Mg^{2+} in neuroblastoma SH-SY5Y cells: a fluorescence and 31P NMR study. *Biophysical Journal* 1999;76:2934–42.
- Autry AE, Adachi M, Nosyreva E, Na ES, Los MF, Cheng PF, et al. NMDA receptor blockade at rest triggers rapid behavioural antidepressant responses. *Nature* 2011;475:91–5.
- Barragan-Rodriguez L, Rodriguez-Moran M, Guerrero-Romero F. Efficacy and safety of oral magnesium supplementation in the treatment of depression in the elderly with type 2 diabetes: a randomized, equivalent trial. *Magnesium Research* 2008;21:218–23.
- Barria A, Malinow R. NMDA receptor subunit composition controls synaptic plasticity by regulating binding to CaMKII. *Neuron* 2005;48:289–301.
- Begon S, Pickering G, Eschaliere A, Mazur A, Rayssiguier Y, Dubray C. Role of spinal NMDA receptors, protein kinase C and nitric oxide synthase in the hyperalgesia induced by magnesium deficiency in rats. *British Journal of Pharmacology* 2001;134:1227–36.
- Beneyto M, Meador-Woodruff JH. Lamina-specific abnormalities of NMDA receptor-associated postsynaptic protein transcripts in the prefrontal cortex in schizophrenia and bipolar disorder. *Neuropsychopharmacology* 2008;33:2175–86.
- van Berckel BN, Oranje B, van Ree JM, Verbaten MN, Kahn RS. The effects of low dose ketamine on sensory gating, neuroendocrine secretion and behavior in healthy human subjects. *Psychopharmacology (Berl)* 1998;137:271–81.
- Berman RM, Cappiello A, Anand A, Oren DA, Heninger GR, Charney DS, et al. Antidepressant effects of ketamine in depressed patients. *Biological Psychiatry* 2000;47:351–4.
- Bettendorff L, Sallanon-Moulin M, Touret M, Wins P, Margineanu I, Schoffeniels E. Paradoxical sleep deprivation increases the content of glutamate and glutamine in rat cerebral cortex. *Sleep* 1996;19:65–71.
- Blair RE, Churn SB, Sombati S, Lou JK, DeLorenzo RJ. Long-lasting decrease in neuronal Ca^{2+} /calmodulin-dependent protein kinase II activity in a hippocampal neuronal culture model of spontaneous recurrent seizures. *Brain Research* 1999;851:54–65.
- Broughton Pipkin F, Waldron BA. Ketamine hypertension and the renin–angiotensin system. *Clinical and Experimental Hypertension A* 1983;5:875–83.
- Burgdorf J, Zhang XL, Nicholson KL, Balster RL, David Leander J, Stanton PK, et al. GLYX-13, an NMDA receptor glycine-site functional partial agonist, induces

- antidepressant-like effects without ketamine-like side effects. *Neuropsychopharmacology* 2013.
- Burnashev N, Monyer H, Seeburg PH, Sakmann B. Divalent ion permeability of AMPA receptor channels is dominated by the edited form of a single subunit. *Neuron* 1992;8:189–98.
- Camardese G, De Risio L, Pizi G, Mattioli B, Buccelletti F, Serrani R, et al. Plasma magnesium levels and treatment outcome in depressed patients. *Nutritional Neuroscience* 2012;15:78–84.
- Carlezon Jr WA, Duman RS, Nestler EJ. The many faces of CREB. *Trends in Neurosciences* 2005;28:436–45.
- Caudal D, Godsil BP, Mailliet F, Bergerot D, Jay TM. Acute stress induces contrasting changes in AMPA receptor subunit phosphorylation within the prefrontal cortex, amygdala and hippocampus. *PLoS One* 2010;5:e15282.
- Chatterjee M, Verma R, Ganguly S, Palit G. Neurochemical and molecular characterization of ketamine-induced experimental psychosis model in mice. *Neuropharmacology* 2012.
- Chen H, Huang LY. Protein kinase C reduces Mg²⁺ block of NMDA-receptor channels as a mechanism of modulation. *Nature* 1992;356:521–3.
- Cheng C, Reynolds JJ. Subcellular localization of glutamate-stimulated intracellular magnesium concentration changes in cultured rat forebrain neurons using confocal microscopy. *Neuroscience* 2000;95:973–9.
- Chiba S, Numakawa T, Ninomiya M, Richards MC, Wakabayashi C, Kunugi H. Chronic restraint stress causes anxiety- and depression-like behaviors, downregulates glucocorticoid receptor expression, and attenuates glutamate release induced by brain-derived neurotrophic factor in the prefrontal cortex. *Progress in Neuro-Psychopharmacology & Biological Psychiatry* 2012;39:112–9.
- Chollet D, Franken P, Raffin Y, Malafosse A, Widmer J, Tafti M. Blood and brain magnesium in inbred mice and their correlation with sleep quality. *American Journal of Physiology – Regulatory, Integrative and Comparative Physiology* 2000;279:R2173–8.
- Choudary PV, Molnar M, Evans SJ, Tomita H, Li JZ, Vawter MP, et al. Altered cortical glutamatergic and GABAergic signal transmission with glial involvement in depression. *Proceedings of the National Academy of Sciences of the United States of America* 2005;102:15653–8.
- Cirelli C, Tononi G. Gene expression in the brain across the sleep-waking cycle. *Brain Research* 2000;885:303–21.
- D'Sa C, Duman RS. Antidepressants and neuroplasticity. *Bipolar Disorders* 2002;4:183–94.
- van Donkelaar EL, van den Hove DL, Blokland A, Steinbusch HW, Prickaerts J. Stress-mediated decreases in brain-derived neurotrophic factor as potential confounding factor for acute tryptophan depletion-induced neurochemical effects. *European Neuropsychopharmacology* 2009;19:812–21.
- Douglas BG, Dagirmanjian R. The effects of magnesium deficiency of ketamine sleeping times in the rat. *British Journal of Anaesthesia* 1975;47:336–40.
- Duman RS. Novel therapeutic approaches beyond the serotonin receptor. *Biological Psychiatry* 1998;44:324–35.
- Duman RS, Li N, Liu RJ, Duric V, Aghajanian G. Signaling pathways underlying the rapid antidepressant actions of ketamine. *Neuropharmacology* 2012;62:35–41.
- Duncan JR WC, Selter J, Brutsche N, Sarasso S, Zarate Jr CA. Baseline delta sleep ratio predicts acute ketamine mood response in major depressive disorder. *Journal of Affective Disorders* 2012.
- Duncan WC, Sarasso S, Ferrarelli F, Selter J, Riedner BA, Hejazi NS, et al. Concomitant BDNF and sleep slow wave changes indicate ketamine-induced plasticity in major depressive disorder. *International Journal of Neuropsychopharmacology* 2012;1–11.
- Dunner DL, Meltzer HL, Schreiner HC, Feigelson JL. Plasma and erythrocyte magnesium levels in patients with primary affective disorder during chronic lithium treatment. *Acta Psychiatrica Scandinavica* 1975;51:104–9.
- Dwivedi Y, Rao JS, Rizavi HS, Kotowski J, Conley RR, Roberts RC, et al. Abnormal expression and functional characteristics of cyclic adenosine monophosphate response element binding protein in postmortem brain of suicide subjects. *Archives of General Psychiatry* 2003;60:273–82.
- Ebel H, Wolff JR, Dorn F, Gunther T. The effect of hormones on the levels of electrolytes, ATPase and the structure of the endoplasmic reticulum in rat brain. *Zeitschrift für Klinische Chemie und Klinische Biochemie* 1971;9:249–56.
- Ebel H, Hollstein M, Gunther T. Differential effect of imipramine and related compounds on Mg²⁺ efflux from rat erythrocytes. *Biochimica et Biophysica Acta* 2004;1667:132–40.
- Eby GA, Eby KL. Rapid recovery from major depression using magnesium treatment. *Medical Hypotheses* 2006.
- Eby GA, Eby KL, Murck H. Magnesium and major depression. In: Vink R, Nechifor M, editors. *Magnesium in the central nervous system*. Adelaide: University of Adelaide Press; 2011. p. 313–30.
- Emanuele E, Geroldi D, Minoretto P, Coen E, Politi P. Increased plasma aldosterone in patients with clinical depression. *Archives of Medical Research* 2005;36:544–8.
- Franklin M, Bermudez I, Hlavacova N, Babic S, Murck H, Schmuckermair C, et al. Aldosterone increases earlier than corticosterone in new animal models of depression: is this an early marker? *Journal of Psychiatric Research* 2012;46:1394–7.
- Feyissa AM, Chandran A, Stockmeier CA, Karolewicz B. Reduced levels of NR2A and NR2B subunits of NMDA receptor and PSD-95 in the prefrontal cortex in major depression. *Progress in Neuro-Psychopharmacology & Biological Psychiatry* 2009;33:70–5.
- Fleischer-Lambropoulos E, Kazazoglou T, Geladopoulos T, Kentroti S, Stefanis C, Vernadakis A. Stimulation of glutamine synthetase activity by excitatory amino acids in astrocyte cultures derived from aged mouse cerebral hemispheres may be associated with non-N-methyl-D-aspartate receptor activation. *International Journal of Developmental Neuroscience* 1996;14:523–30.
- Fortin DA, Davare MA, Srivastava T, Brady JD, Nygaard S, Derkach VA, et al. Long-term potentiation-dependent spine enlargement requires synaptic Ca²⁺-permeable AMPA receptors recruited by CaM-kinase I. *Journal of Neuroscience* 2010;30:11565–75.
- Fortin DA, Srivastava T, Dwarakanath D, Pierre P, Nygaard S, Derkach VA, et al. Brain-derived neurotrophic factor activation of CaM-kinase kinase via transient receptor potential canonical channels induces the translation and synaptic incorporation of GluA1-containing calcium-permeable AMPA receptors. *Journal of Neuroscience* 2012;32:8127–37.
- Guzman-Marin R, Ying Z, Sunstova N, Methippara M, Bashir T, Szymusiak R, et al. Suppression of hippocampal plasticity-related gene expression by sleep deprivation in rats. *Journal of Physiology* 2006;575:807–19.
- Gomez-Villafuertes R, del Puerto A, Diaz-Hernandez M, Bustillo D, Diaz-Hernandez JI, Huerta PG, et al. Ca²⁺/calmodulin-dependent kinase II signalling cascade mediates P2X7 receptor-dependent inhibition of neurogenesis in neuroblastoma cells. *FEBS Journal* 2009;276:5307–25.
- Gorelik O, Efrati S, Berman S, Almozni-Saraifan D, Alon I, Shteinshnaider M, et al. Effect of various clinical variables on total intracellular magnesium in hospitalized normomagnesemic diabetic patients before discharge. *Biological Trace Element Research* 2007;120:102–9.
- Gourley SL, Wu FJ, Kiraly DD, Ploski JE, Kedves AT, Duman RS, et al. Regionally specific regulation of ERK MAP kinase in a model of antidepressant-sensitive chronic depression. *Biological Psychiatry* 2008;63:353–9.
- Green EK, Grozeva D, Raybould R, Elvidge G, Macgregor S, Craig I, et al. P2RX7: a bipolar and unipolar disorder candidate susceptibility gene? *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics* 2009;150B:1063–9.
- Greenberg J, Lichtenstein N. Effect of manganous and magnesium ions concentration on glutamine synthetase and glutamotransferase of sheep brain. *Journal of Biological Chemistry* 1959;234:2337–9.
- Held K, Antonijevic IA, Kunzel H, Uhr M, Wetter TC, Golly IC, et al. Oral Mg(2+) supplementation reverses age-related neuroendocrine and sleep EEG changes in humans. *Pharmacopsychiatry* 2002;35:135–43.
- Hemant Kumar BS, Mishra SK, Rana P, Singh S, Khushu S. Neurodegenerative evidences during early onset of depression in CMS rats as detected by proton magnetic resonance spectroscopy at 7 T. *Behavioural Brain Research* 2012;232:53–9.
- Hemmeter UM, Hemmeter-Spernal J, Krieg JC. Sleep deprivation in depression. *Expert Review of Neurotherapeutics* 2010;10:1101–15.
- Hlavacova N, Wes PD, Ondrejckova M, Flynn ME, Poundstone PK, Babic S, et al. Subchronic treatment with aldosterone induces depression-like behaviours and gene expression changes relevant to major depressive disorder. *International Journal of Neuropsychopharmacology* 2011;1–19.
- Hoyt KR, Rajdev S, Fattman CL, Reynolds JJ. Cyclothiazide modulates AMPA receptor-mediated increases in intracellular free Ca²⁺ and Mg²⁺ in cultured neurons from rat brain. *Journal of Neurochemistry* 1995;64:2049–56.
- Huang CY, Liou YF, Chung SY, Lin WY, Jong GP, Kuo CH, et al. Role of ERK signaling in the neuroprotective efficacy of magnesium sulfate treatment during focal cerebral ischemia in the gerbil cortex. *Chinese Journal of Physiology* 2010;53:299–309.
- Iasevoli F, Polese D, Ambesi-Impiombato A, Muscettola G, de Bartolomeis A. Ketamine-related expression of glutamatergic postsynaptic density genes: possible implications in psychosis. *Neuroscience Letters* 2007;416:1–5.
- Iosifescu DV, Bolo NR, Nierenberg AA, Jensen JE, Fava M, Renshaw PF. Brain bioenergetics and response to triiodothyronine augmentation in major depressive disorder. *Biological Psychiatry* 2008;63:1127–34.
- Jiang LH. Inhibition of P2X(7) receptors by divalent cations: old action and new insight. *European Biophysics Journal* 2009;38:339–46.
- Jiang Q, Wang J, Wu X, Jiang Y. Alterations of NR2B and PSD-95 expression after early-life epileptiform discharges in developing neurons. *International Journal of Developmental Neuroscience* 2007;25:165–70.
- Jiang Q, Wang J, Wu Y, Wu X, Qin J, Jiang Y. Early-life epileptiform discharges exert both rapid and long-lasting effects on AMPAR subunit composition and distribution in developing neurons. *Neuroscience Letters* 2008;444:31–5.
- Kalkman HO. Circumstantial evidence for a role of glutamine-synthetase in suicide. *Medical Hypotheses* 2011;76:905–7.
- Karolewicz B, Szebeni K, Gilmore T, Maciag D, Stockmeier CA, Ordway GA. Elevated levels of NR2A and PSD-95 in the lateral amygdala in depression. *International Journal of Neuropsychopharmacology* 2009;12:143–53.
- Khine K, Rosenstein DL, Elin RJ, Niemela JE, Schmidt PJ, Rubinow DR. Magnesium (mg) retention and mood effects after intravenous mg infusion in premenstrual dysphoric disorder. *Biological Psychiatry* 2006;59:327–33.
- Kielczykowska M, Pasternak K, Musik I. The influence of lithium on calcium and magnesium homeostasis in serum and tissues of rats. *Annales Universitatis Mariae Curie-Skłodowska* 2003;58:281–4.
- Kielczykowska M, Musik I, Hordyjewska A, Boguszewska A, Lewandowska A, Pasternak K. Oral administration of lithium increases tissue magnesium contents but not plasma magnesium level in rats. *Pharmacological Reports* 2007;59:291–5.
- Kim SJ, Kang HS, Lee MY, Lee SJ, Seol JW, Park SY, et al. Ketamine-induced cardiac depression is associated with increase in [Mg²⁺]_i and activation of p38 MAP

- kinase and ERK 1/2 in guinea pig. *Biochemical and Biophysical Research* 2006;349:716–22.
- King LJ, Carl JL, Archer EG, Castellonet M. Effects of lithium on brain energy reserves and cations in vivo. *Journal of Pharmacology and Experimental Therapeutics* 1969;168:163–70.
- Knox D, Perrine SA, George SA, Galloway MP, Liberzon I. Single prolonged stress decreases glutamate, glutamine, and creatine concentrations in the rat medial prefrontal cortex. *Neuroscience Letters* 2010;480:16–20.
- Kuipers SD, Trentani A, Den Boer JA, Ter Horst GJ. Molecular correlates of impaired prefrontal plasticity in response to chronic stress. *Journal of Neurochemistry* 2003;85:1312–23.
- Kuner T, Schoepfer R. Multiple structural elements determine subunit specificity of Mg²⁺ block in NMDA receptor channels. *Journal of Neuroscience* 1996;16:3549–58.
- Kunzel HE, Apostolopoulou K, Pallauf A, Gerum S, Merkle K, Schulz S, et al. Quality of life in patients with primary aldosteronism: Gender differences in untreated and long-term treated patients and associations with treatment and aldosterone. *Journal of Psychiatric Research* 2012;46:1650–4.
- Laifenfeld D, Karry R, Grauer E, Klein E, Ben-Shachar D. Antidepressants and prolonged stress in rats modulate CAM-L1, laminin, and pCREB, implicated in neuronal plasticity. *Neurobiology of Disease* 2005;20:432–41.
- Landfield PW, Morgan GA. Chronically elevating plasma Mg²⁺ improves hippocampal frequency potentiation and reversal learning in aged and young rats. *Brain Research* 1984;322:167–71.
- Lante F, Toledo-Salas JC, Ondrejčák T, Rowan MJ, Ulrich D. Removal of synaptic Ca²⁺-permeable AMPA receptors during sleep. *Journal of Neuroscience* 2011;31:3953–61.
- Lee YA, Goto Y. Chronic stress modulation of prefrontal cortical NMDA receptor expression disrupts limbic structure-prefrontal cortex interaction. *European Journal of Neuroscience* 2011;34:426–36.
- Lee HK, Barbarosie M, Kameyama K, Bear MF, Huganir RL. Regulation of distinct AMPA receptor phosphorylation sites during bidirectional synaptic plasticity. *Nature* 2000;405:955–9.
- Lee MY, Kim SJ, Kim JS, Kang HS. Imipramine-induced cardiac depression is responsible for the increase in intracellular magnesium and the activation of ERK 1/2 in rats. *Journal of Cardiovascular Pharmacology and Therapeutics* 2010;15:303–10.
- Lee M, Jantarantotai N, McGeer E, McLarnon JG, McGeer PL. Mg²⁺ ions reduce microglial and THP-1 cell neurotoxicity by inhibiting Ca²⁺ entry through purinergic channels. *Brain Research* 2011;1369:21–35.
- Leon D, Hervas C, Miras-Portugal MT. P2Y1 and P2X7 receptors induce calcium/calmodulin-dependent protein kinase II phosphorylation in cerebellar granule neurons. *European Journal of Neuroscience* 2006;23:2999–3013.
- Li N, Lee B, Liu RJ, Banasr M, Dwyer JM, Iwata M, et al. mTOR-dependent synapse formation underlies the rapid antidepressant effects of NMDA antagonists. *Science* 2010;329:959–64.
- Li N, Liu RJ, Dwyer JM, Banasr M, Lee B, Son H, et al. Glutamate N-methyl-D-aspartate receptor antagonists rapidly reverse behavioral and synaptic deficits caused by chronic stress exposure. *Biological Psychiatry* 2011a;69:754–61.
- Li X, Frye MA, Shelton RC. Review of pharmacological treatment in mood disorders and future directions for drug development. *Neuropsychopharmacology* 2011.
- Li-Smerin Y, Aizenman E, Johnson JW. Inhibition by intracellular Mg²⁺ of recombinant N-methyl-D-aspartate receptors expressed in Chinese hamster ovary cells. *Journal of Pharmacology and Experimental Therapeutics* 2000;292:1104–10.
- Lisman J, Yasuda R, Raghavachari S. Mechanisms of CaMKII action in long-term potentiation. *Nature Reviews Neuroscience* 2012;13:169–82.
- Liu HT, Hollmann MW, Liu WH, Hoenemann CW, Durieux ME. Modulation of NMDA receptor function by ketamine and magnesium: part I. *Anesthesia & Analgesia* 2001;92:1173–81.
- Loix S, De Kock M, Henin P. The anti-inflammatory effects of ketamine: state of the art. *Acta Anaesthesiologica Belgica* 2011;62:47–58.
- Lucae S, Salyakina D, Barden N, Harvey M, Gagne B, Labbe M, et al. P2RX7, a gene coding for a purinergic ligand-gated ion channel, is associated with major depressive disorder. *Human Molecular Genetics* 2006;15:2438–45.
- Magistretti PJ. Role of glutamate in neuron-glia metabolic coupling. *American Journal of Clinical Nutrition* 2009;90:875S–80S.
- Manji HK, Duman RS. Impairments of neuroplasticity and cellular resilience in severe mood disorders: implications for the development of novel therapeutics. *Psychopharmacology Bulletin* 2001;35:5–49.
- Maurizi MR, Pinkofsky HB, McFarland PJ, Ginsburg A. Mg²⁺ is bound to glutamine synthetase extracted from bovine or ovine brain in the presence of L-methionine-S-sulfoximine phosphate. *Archives of Biochemistry and Biophysics* 1986;246:494–500.
- Maurizi MR, Pinkofsky HB, Ginsburg A. ADP, chloride ion, and metal ion binding to bovine brain glutamine synthetase. *Biochemistry* 1987;26:5023–31.
- Maret S, Faraguna U, Nelson AB, Cirelli C, Tononi G. Sleep and waking modulate spine turnover in the adolescent mouse cortex. *Nature Neuroscience* 2011;14:1418–20.
- McIntosh TK, Vink R, Soares H, Hayes R, Simon R. Effect of noncompetitive blockade of N-methyl-D-aspartate receptors on the neurochemical sequelae of experimental brain injury. *Journal of Neurochemistry* 1990;55:1170–9.
- Muller A, Gunzel D, Schlue WR. Activation of AMPA/kainate receptors but not acetylcholine receptors causes Mg²⁺ influx into Retzius neurons of the leech *Hirudo medicinalis*. *Journal of General Physiology* 2003;122:727–39.
- Murck H. Magnesium and affective disorders. *Nutritional Neuroscience* 2002;5:375–89.
- Murck H, Held K, Ziegenbein M, Kunzel H, Koch K, Steiger A. The renin-angiotensin-aldosterone system in patients with depression compared to controls—a sleep endocrine study. *BMC Psychiatry* 2003;3:15.
- Murck H, Schubert MI, Schmid D, Schussler P, Steiger A, Auer DP. The glutamatergic system and its relation to the clinical effect of therapeutic-sleep deprivation in depression — an MR spectroscopy study. *Journal of Psychiatric Research* 2009;43:175–80.
- Murck H, Schussler P, Steiger A. Renin-angiotensin-aldosterone system: the forgotten stress hormone system: relationship to depression and sleep. *Pharmacopsychiatry* 2012.
- Nechifor M. Interactions between magnesium and psychotropic drugs. *Magnesium Research* 2008;21:97–100.
- Orser B, Smith D, Henderson S, Gelb A. Magnesium deficiency increases ketamine sensitivity in rats. *Canadian Journal of Anaesthesia* 1997;44:883–90.
- Park HJ, Kang WS, Paik JW, Kim JW. Effect of valproic acid through regulation of NMDA receptor-ERK signaling in sleep deprivation rats. *Journal of Molecular Neuroscience* 2012;47:554–8.
- Paul IA, Nowak G, Lauer RT, Popik P, Skolnick P. Adaptation of the N-methyl-D-aspartate receptor complex following chronic antidepressant treatments. *Journal of Pharmacology and Experimental Therapeutics* 1994;269:95–102.
- Perraud AL, Zhao X, Ryazanov AG, Schmitz C. The channel-kinase TRPM7 regulates phosphorylation of the translational factor eEF2 via eEF2-k. *Cell Signal* 2011;23:586–93.
- Pittaluga A, Bonfanti A, Raiteri M. Somatostatin potentiates NMDA receptor function via activation of InsP(3) receptors and PKC leading to removal of the Mg²⁺ block without depolarization. *British Journal of Pharmacology* 2000;130:557–66.
- Poleszak E, Wlaz P, Szewczyk B, Kedzierska E, Wyska E, Librowski T, et al. Enhancement of antidepressant-like activity by joint administration of imipramine and magnesium in the forced swim test: behavioral and pharmacokinetic studies in mice. *Pharmacology Biochemistry and Behavior* 2005;81:524–9.
- Poleszak E, Wlaz P, Kedzierska E, Nieoczym D, Wyska E, Szymura-Oleksiak J, et al. Immobility stress induces depression-like behavior in the forced swim test in mice: effect of magnesium and imipramine. *Pharmacological Reports* 2006;58:746–52.
- Poleszak E, Wlaz P, Kedzierska E, Nieoczym D, Wrobel A, Fidecka S, et al. NMDA/glutamate mechanism of antidepressant-like action of magnesium in forced swim test in mice. *Pharmacology Biochemistry and Behavior* 2007;88:158–64.
- Popoli M, Yan Z, McEwen BS, Sanacora G. The stressed synapse: the impact of stress and glucocorticoids on glutamate transmission. *Nature Reviews Neuroscience* 2012;13:22–37.
- Preskorn SH, Baker B, Kolluri S, Menniti FS, Krams M, Landen JW. An innovative design to establish proof of concept of the antidepressant effects of the NR2B subunit selective N-methyl-D-aspartate antagonist, CP-101,606, in patients with treatment-refractory major depressive disorder. *Journal of Clinical Psychopharmacology* 2008;28:631–7.
- Quan MN, Zhang N, Wang YY, Zhang T, Yang Z. Possible antidepressant effects and mechanisms of memantine in behaviors and synaptic plasticity of a depression rat model. *Neuroscience* 2011;182:88–97.
- Rasgon NL, Kenna HA, Williams KE, Powers B, Wroolie T, Schatzberg AF. Rosiglitazone add-on in treatment of depressed patients with insulin resistance: a pilot study. *Scientific World Journal* 2010;10:321–8.
- Reus GZ, Stringari RB, Ribeiro KF, Ferraro AK, Vitto MF, Cesconetto P, et al. Ketamine plus imipramine treatment induces antidepressant-like behavior and increases CREB and BDNF protein levels and PKA and PKC phosphorylation in rat brain. *Behavioural Brain Research* 2011;221:166–71.
- Richards CD, Sercombe R. Calcium, magnesium and the electrical activity of guinea-pig olfactory cox in vitro. *Journal of Physiology* 1970;211:571–84.
- Rothe F, Wolf G, Fischer S, Hass P, Keilhoff G, Abicht K. Quinolate and kainate facilitate magnesium penetration into brain tissue. *Neuroreport* 1993;4:205–7.
- Rowland LM, Bustillo JR, Mullins PG, Jung RE, Lenroot R, Landgraf E, et al. Effects of ketamine on anterior cingulate glutamate metabolism in healthy humans: a 4-T proton MRS study. *American Journal of Psychiatry* 2005;162:394–6.
- Rubin RR, Knowler WC, Ma Y, Marrero DG, Edelstein SL, Walker EA, et al. Depression symptoms and antidepressant medicine use in diabetes prevention program participants. *Diabetes Care* 2005;28:830–7.
- Runyan AL, Sun Y, Bhattacharya SK, Ahokas RA, Chhokar VS, Gerling IC, et al. Responses in extracellular and intracellular calcium and magnesium in aldosteronism. *Journal of Laboratory and Clinical Medicine* 2005;146:76–84.
- Salehi-Sadaghiani M, Javadi-Paydar M, Gharedaghi MH, Zandieh A, Heydarpour P, Yousefzadeh-Fard Y, et al. NMDA receptor involvement in antidepressant-like effect of pioglitazone in the forced swimming test in mice. *Psychopharmacology (Berl)* 2012.
- Salvadore G, van der Veen JW, Zhang Y, Marenco S, Machado-Vieira R, Baumann J, et al. An investigation of amino-acid neurotransmitters as potential predictors of clinical improvement to ketamine in depression. *International Journal of Neuropsychopharmacology* 2011;1–10.
- Sanacora G, Kendell SF, Fenton L, Coric V, Krystal JH. Riluzole augmentation for treatment-resistant depression. *American Journal of Psychiatry* 2004;161:2132.

- Sanacora G, Kendell SF, Levin Y, Simen AA, Fenton LR, Coric V, et al. Preliminary evidence of riluzole efficacy in antidepressant-treated patients with residual depressive symptoms. *Biological Psychiatry* 2007;61:822–5.
- Sartori SB, Whittle N, Hetzenauer A, Singewald N. Magnesium deficiency induces anxiety and HPA axis dysregulation: modulation by therapeutic drug treatment. *Neuropharmacology* 2012;62:304–12.
- Sepanjnia K, Modabbernia A, Ashrafi M, Modabbernia MJ, Akhondzadeh S. Pioglitazone adjunctive therapy for moderate-to-severe major depressive disorder: randomized double-blind placebo-controlled trial. *Neuropsychopharmacology* 2012;37:2093–100.
- Shapira Y, Lam AM, Artru AA, Eng C, Soltow L. Ketamine alters calcium and magnesium in brain tissue following experimental head trauma in rats. *Journal of Cerebral Blood Flow & Metabolism* 1993;13:962–8.
- Shu L, Li T, Han S, Ji F, Pan C, Zhang B, et al. Inhibition of neuron-specific CREB dephosphorylation is involved in propofol and ketamine-induced neuroprotection against cerebral ischemic injuries of mice. *Neurochemical Research* 2012;37:49–58.
- Silver BB. Development of cellular magnesium nano-analysis in treatment of clinical magnesium deficiency. *Journal of the American College of Nutrition* 2004;23:732S–7S.
- Singewald N, Sinner C, Hetzenauer A, Sartori SB, Murck H. Magnesium-deficient diet alters depression- and anxiety-related behavior in mice – influence of desipramine and *Hypericum perforatum* extract. *Neuropharmacology* 2004;47:1189–97.
- Skaper SD, Debetto P, Giusti P. The P2X7 purinergic receptor: from physiology to neurological disorders. *FASEB Journal* 2010;24:337–45.
- Slutsky I, Abumaria N, Wu LJ, Huang C, Zhang L, Li B, et al. Enhancement of learning and memory by elevating brain magnesium. *Neuron* 2010;65:165–77.
- Soronen P, Mantere O, Melartin T, Suominen K, Vuorilehto M, Rytasala H, et al. P2RX7 gene is associated consistently with mood disorders and predicts clinical outcome in three clinical cohorts. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics* 2011;156B:435–47.
- Sun L, June Liu S. Activation of extrasynaptic NMDA receptors induces a PKC-dependent switch in AMPA receptor subtypes in mouse cerebellar stellate cells. *Journal of Physiology* 2007;583:537–53.
- Tanis KQ, Duman RS, Newton SS. CREB binding and activity in brain: regional specificity and induction by electroconvulsive seizure. *Biological Psychiatry* 2008;63:710–20.
- Taylor MJ, Tiangga ER, Mhuirchearthaigh RN, Cowen PJ. Lack of effect of ketamine on cortical glutamate and glutamine in healthy volunteers: a proton magnetic resonance spectroscopy study. *Journal of Psychopharmacology* 2012;26:733–7.
- Teyssier JR, Ragot S, Chauvet-Gelinier JC, Trojak B, Bonin B. Activation of a Delta-FOSB dependent gene expression pattern in the dorsolateral prefrontal cortex of patients with major depressive disorder. *Journal of Affective Disorders* 2011;133:174–8.
- Trullas R, Skolnick P. Functional antagonists at the NMDA receptor complex exhibit antidepressant actions. *European Journal of Pharmacology* 1990;185:1–10.
- Vanhoutte P, Bading H. Opposing roles of synaptic and extrasynaptic NMDA receptors in neuronal calcium signalling and BDNF gene regulation. *Current Opinion in Neurobiology* 2003;13:366–71.
- Vyazovskiy VV, Cirelli C, Pfister-Genskow M, Faraguna U, Tononi G. Molecular and electrophysiological evidence for net synaptic potentiation in wake and depression in sleep. *Nature Neuroscience* 2008;11:200–8.
- Weglicki WB. Hypomagnesemia and inflammation: clinical and basic aspects. *Annual Review of Nutrition* 2012;32:55–71.
- Weglicki WB, Phillips TM, Freedman AM, Cassidy MM, Dickens BF. Magnesium-deficiency elevates circulating levels of inflammatory cytokines and endothelin. *Molecular and Cellular Biochemistry* 1992;110:169–73.
- Wu TC, Chen HT, Chang HY, Yang CY, Hsiao MC, Cheng ML, et al. Mineralocorticoid receptor antagonist spironolactone prevents chronic corticosterone induced depression-like behavior. *Psychoneuroendocrinology* 2012.
- Xing G, Russell S, Hough C, O'Grady J, Zhang L, Yang S, et al. Decreased prefrontal CaMKII alpha mRNA in bipolar illness. *Neuroreport* 2002;13:501–5.
- Xiong ZG, Raouf R, Lu WY, Wang LY, Orser BA, Dudek EM, et al. Regulation of N-methyl-D-aspartate receptor function by constitutively active protein kinase C. *Molecular Pharmacology* 1998;54:1055–63.
- Xiong H, Futamura T, Jourdi H, Zhou H, Takei N, Diverse-Pierluissi M, et al. Neurotrophins induce BDNF expression through the glutamate receptor pathway in neocortical neurons. *Neuropharmacology* 2002;42:903–12.
- Yamada S, Yamamoto M, Ozawa H, Riederer P, Saito T. Reduced phosphorylation of cyclic AMP-responsive element binding protein in the postmortem orbitofrontal cortex of patients with major depressive disorder. *Journal of Neural Transmission* 2003;110:671–80.
- Yu H, Chen ZY. The role of BDNF in depression on the basis of its location in the neural circuitry. *Acta Pharmacologica Sinica* 2011;32:3–11.
- Yuen EY, Wei J, Liu W, Zhong P, Li X, Yan Z. Repeated stress causes cognitive impairment by suppressing glutamate receptor expression and function in prefrontal cortex. *Neuron* 2012;73:962–77.
- Zarate Jr CA, Singh JB, Carlson PJ, Brutsche NE, Ameli R, Luckenbaugh DA, et al. A randomized trial of an N-methyl-D-aspartate antagonist in treatment-resistant major depression. *Archives of General Psychiatry* 2006;63:856–64.
- Zarate Jr CA, Mathews DC, Furey ML. Human Biomarkers of Rapid Antidepressant Effects. *Biological Psychiatry* 2013. Epub ahead of print.
- Zeng J, Xia S, Zhong W, Li J, Lin L. In vitro and in vivo effects of ketamine on generation and function of dendritic cells. *Journal of Pharmacological Sciences* 2011;117:170–9.
- Zhang Y, Zu X, Luo W, Yang H, Luo G, Zhang M, et al. Social isolation produces anxiety-like behaviors and changes PSD-95 levels in the forebrain. *Neuroscience Letters* 2012;514:27–30.